

A case report: 1-year follow-up of giant cystic lymphangioma of right mesocolon

Nguyen Anh Tuan, Nguyen Van Du, Nguyen To Hoai, Pham Van Hiep

Department of Gastrointestinal tract Surgery, 108 Military Central Hospital

Corresponding Author:

Nguyen Van Du
108 Military Central Hospital
No 1, Tran Hung Dao, Hai Ba Trung,
Ha Noi
Mobile: +84975454008
Email: nvdu3108@gmail.com

Received date: 01/8/2022

Accepted date: 30/8/2022

Published date: 15/9/2022

Abstract

Introduction: Cystic lymphangiomas are rare benign tumors of the lymph vessels and are usually found in children. However, abdominal cystic lymphangioma in mesocolon is extremely rare in adult patients.

Case presentation: We reported a 15-year-old female with giant cystic lymphangioma of the right mesocolon. On examination, only abdominal pain was confirmed. Abdominal computed tomography (CT) showed a large multiseptated cystic mass. The patient underwent a total right mesocolic excision with the lesion. The patient recovered well on postoperative follow-up and was discharged on the fifth day. No evidence of recurrence had also been found during 1-year follow-up period.

Conclusion: Complete tumor removal is the optimal choice for the management of intra-abdominal cystic lymphangioma. However, incomplete resection may lead to local recurrence.

Keywords: Cystic, Lymphangioma, Mesocolon.

Introduction

Cystic lymphangioma is a benign tumor of the lymphatic system, mainly affecting the craniofacial region, neck, or chest, common in childhood. Cystic lymphangioma is extremely rare in the adult population, with about 1 in 250,000 hospital admission [1], [2]. In the intraperitoneal cavity, cystic lymphangioma was found in 70% of the small mesenteric intestine and the other in the greater omentum, mesocolon, and retroperitoneum [3]. The diagnosis is still challenging due to confusion with mesenteric cysts. We describe a case of giant cystic lymphangioma of mesocolon that was successfully performed tumor removal. This case report has been

reported in line with the SCARE criteria [4].

Case presentation

15-year-old female (BMI: 16.6 kg/m²) presented to our hospital with 3 months of abdominal pain. She denied vomiting, nausea, and a history of any other abdominal disorder. Physical examination revealed a palpable tumor from the epigastric to hypogastric region, especially on the right side of the umbilicus. There were unremarkable in routine laboratory and hematologic investigations. Computed tomography showed a large cystic mass with 16.0 x 18.2 cm in size, extending from the lower of the liver to the ileocecal corner, suspecting of a mesenteric cyst or omental lymphangioma (Figure 1).



Figure 1: Computed tomography showed a large cystic mass.

During exploration surgery, a huge cystic mass was found covering the right mesocolon (Figure 2). The surface of the mass showed fibrotic capsules with thick walls. Because of the inability to separate the descending colon and mesocolon from the lesion, total right mesocolic excision with the mass was

performed simultaneously. A side-to-side anastomosis was constructed between the terminal ileum and the transverse colon using a linear cutter stapler. After resection, the cystic tumor was grossly measured at 15.2 x 16.8 x 3.2 cm (Figure 3). The operative time was 254 minutes, and the blood loss was 350ml.



Figure 2: Intraoperative view remove of a giant cystic lymphangioma



Figure 3: Gross specimen showed a cystic lymphangioma with multiseptated.




Figure 4: Histology of lymphangioma. The cystic wall consisted of fibroconnective tissue accompanied by dilated lymphatic spaces and lymphoid cell aggregations in the endothelial lining of lymphatic vessels (A) H&E × 40, (B) H&E × 200.

Histologically, the cystic wall consisted of fibrous tissue, lymphatic tissue, blood vessels, and smooth muscle, with an inner surface of a single layer of squamous or cuboidal epithelium and no sign of malignancy (Figure 4). Therefore, the final

diagnosis was cystic lymphangioma.

The patient recovered well on postoperative follow-up and was discharged on the fifth day. No evidence of recurrence was also found during 1-year follow-up (Figure 5).




Figure 5: Computed tomography at 1-year follow-up: no recurrence was recognized

Discussion

Cystic lymphangiomas are rare benign tumors that are found predominantly in children. They generally occur in the neck (75%) and the axillary region (20%). Intra-abdominal lymphoma comprises less than 1%, with the most common location in the small bowel mesentery (70%) and extremely rare in the mesocolon [3], [5].

The cause of lymphangioma is still unknown. However, the mechanism of inflammation and fibrosis was accepted by most researchers. No malignant transformation of lymphangioma has been introduced in the literature [6], [7].

The clinical characteristics of intra-abdominal lymphangiomas are variable. Abdominal pain and a palpable mass are the common symptoms. Several complications can cause the acute condition of the patient, including a hemorrhaging cyst, a secondary infection, obstruction, or anemia [8, 9]. Spontaneous rupture of cysts, gastrointestinal bleeding, and cystic peritoneal lymphangiomatosis is rare presented [9], [10].

The diagnosis of intra-abdominal cystic lymphoma is quite difficult because the clinical symptoms are nonspecific. Allen reported a rate of 40% asymptomatic patients in his study and

was discovered incidentally on an imaging exam. Even when the patient underwent an ultrasound and CT scan of the abdomen, confusion could still occur because the images are relatively similar to the mesenteric and omental cysts [2]. Abdominal magnetic resonance imaging (MRI) can show better clarification of the nature of the cystic contents. However, it is unable to determine the origin of the tumor [11].

For non-surgical treatment, some sclerosing therapies using Bleomycin, Tissuecol, OK-432, and Ethibloc were reported as initial treatments of cystic lymphangiomas. These can reduce the size of the mass. Consequently, these can minimize the frequency and extent of surgical intervention [12], [13], [14]. However, Alqahtani et al. have demonstrated that these therapies may cause long-term consequences, such as local recurrences with a very high proportion, up to 100% [15]. Therefore, the optimal choice for intra-abdominal lymphangiomas management is surgery to remove tumors. In most cases of small cysts, total excision without affecting other organs is a priority.

For tumors in solid organs including the spleen, pancreas, liver, or adrenal gland, resection of the lesion along with involved parenchyma should be considered. For cystic lymphangiomas of the mesentery (mesocolon and mesentery), total surgical resection of the tumor with or without bowel can be performed [16]. In order to avoid complications of lymphatic leakage following surgery, a careful examination of the intraoperative lymphatic system is necessary. In this case, we used Ligasure to seal lymphatic ducts and small vessels. After incomplete resection, a recurrence rate of up to 40% was reported. In microscopic residue, the recurrence rate ranges from 5% to 10% [15], [16].

Conclusion

Intra-abdominal cystic lymphangiomas are rare benign tumors that present a diagnostic challenge of diagnosis. Accurate diagnosis is usually based on histopathology. Abdominal ultrasound and CT

scan show a fairly clear cystic mass. Complete resection of the tumor is the optimal choice for lymphangioma management.

References

1. Kumar S, Agrawal N, Khanna R, Khanna AK. Giant lymphatic cyst of omentum: a case report. *Cases J.* 2009;2:23-.
2. Allen JG, Riial TS, Cameron JL, Askin FB, Hruban RH, Campbell KA. Abdominal lymphangiomas in adults. *Journal of Gastrointestinal Surgery.* 2006;10(5):746-51.
3. Kim JH, Ryu WS, Min BW, Song TJ, Son GS, Kim SJ, et al. Acquired omental cystic lymphangioma after subtotal gastrectomy: a case report. *J Korean Med Sci.* 2009;24(6):1212-5.
4. Agha RA, Franchi T, Sohrabi C, Mathew G, Kerwan A. The SCARE 2020 Guideline: Updating Consensus Surgical CAse REport (SCARE) Guidelines. *International journal of surgery.* 2020;84:226-30.
5. Limdi JK, Mehdi S, Sapundzieski M, Manu M, Abbasi AM. Cystic Lymphangioma of the Mesocolon. *Journal of Gastrointestinal Surgery.* 2010;14(9):1459-61.
6. Hornick JL, Fletcher CD. Intraabdominal cystic lymphangiomas obscured by marked superimposed reactive changes: clinicopathological analysis of a series. *Human pathology.* 2005;36(4):426-32.
7. Tsukamoto T, Tanaka S, Yamamoto T, Kakinoki E, Uemichi A, Kubo S, et al. Laparoscopic Excision of a Retroperitoneal Cystic Lymphangioma: Report of a Case. *Surgery Today.* 2003;33(2):142-4.
8. Akashige T, Sato K, Odajima H, Yamazaki S. A case report of recurrent intussusception caused by small bowel lymphangioma in an adult. *International Journal of Surgery Case Reports.* 2020;75:126-30.
9. Lepre L, Costa G, Baldini D, Cortese F, Saputelli A, Gioffre A, et al. Emergency presentation of cystic lymphangioma of the colon: A case report and literature review. *International Journal of Surgery Case Reports.* 2016;24:162-5.
10. Ha TK, Paik SS, Lee KG. Cystic lymphangioma arising from mesocolon. *Clinical gastroenterology and hepatology : the official clinical practice journal of the American Gastroenterological Association.* 2009;7(3):e14-5.
11. Kitamura H, Yamamoto D, Kadoya S, Bando H, Okayama Y, Minato H. Jejunal mesenteric lymphangioma treated by pancreaticoduodenectomy: A case report. *International Journal of Surgery Case Reports.* 2020;77:165-9.

12. Hall N, Ade-Ajayi N, Brewis C, Roebuck DJ, Kiely EM, Drake DP, et al. Is intralesional injection of OK-432 effective in the treatment of lymphangioma in children? *Surgery*. 2003;133(3):238-42.
13. Okada A, Kubota A, Fukuzawa M, Imura K, Kamata S. Injection of bleomycin as a primary therapy of cystic lymphangioma. *Journal of pediatric surgery*. 1992;27(4):440-3.
14. Martinot V, Descamps S, Février P, Patenotre P, Brevière JM, Piette F, et al. [Evaluation of the treatment of cystic lymphangioma by percutaneous injection of Ethibloc in 20 patients]. *Archives de pediatrie : organe officiel de la Societe francaise de pediatrie*. 1997;4(1):8-14.
15. Alqahtani A, Nguyen LT, Flageole H, Shaw K, Laberge JM. 25 years' experience with lymphangiomas in children. *Journal of pediatric surgery*. 1999;34(7):1164-8.
16. Makni A, Chebbi F, Fetirich F, Ksantini R, Bedioui H, Jouini M, et al. Surgical Management of Intra-Abdominal Cystic Lymphangioma. Report of 20 Cases. *World Journal of Surgery*. 2012;36(5):1037-43.